

ENVIRONMENTAL STATEMENT - VOLUME 3 - APPENDIX 6.5

Operation Phase Air Quality Assessment Results Tables: Ecological Receptors

Drax Bioenergy with Carbon Capture and Storage

The Infrastructure Planning (Applications: Prescribed Forms and Procedure) Regulations 2009 - 5(2)(a)

Document Reference Number: 6.3.6.5

Applicant: Drax Power Limited **PINS Reference:** EN010120



REVISION: 0504

DATE: June July 2023

DOCUMENT OWNER: WSP UK Limited

AUTHOR: B Tuckett-Jones APPROVER: N. Ashworth

PUBLIC

TABLE OF CONTENTS

1. OPERATION PHASE AIR QUALITY ASSESSMENT RESULTS TABLES: ECOLOGICAL RECEPTORS
1.1. Impacts on Ecological Receptors1
TABLES
Table 1.1 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NO _x
Table 1.2 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NO _x
Table 1.3 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH ₃ 5
Table 1.4 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean SO ₂ 6
Table 1.5 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate
Table 1.6 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Acid Deposition Rate
Table 1.7 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NO _x (Including Mitigation)12
Table 1.8 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NO _x (Including Mitigation)
Table 1.9 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH ₃ (Including Mitigation)14
Table 1.10 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean SO ₂ (Including Mitigation)
Table 1.11 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate (Including Mitigation)
Table 1.12 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Acid Deposition Rate (Including Mitigation)
Table 1.13 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean NO _x
Table 1.14 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Daily Mean NO _x

able 1.15 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean IH₃22
able 1.16 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean
able 1.17 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate25
Table 1.18 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Acid Deposition Rate
able 1.19 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean IOx (Worst Case Emissions Profile)
able 1.20 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean Ox (Worst Case Emissions Profile)
able 1.21 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean IH3 (Worst Case Emissions Profile)
Table 1.22 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean SO ₂ (Worst Case Emissions Profile)31
able 1.23 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual litrogen Deposition Rate (Worst Case Emissions Profile)32
Table 1.24 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Acid Deposition Rate (Worst Case Emissions Profile)

1. OPERATION PHASE AIR QUALITY ASSESSMENT RESULTS TABLES: ECOLOGICAL RECEPTORS

1.1. IMPACTS ON ECOLOGICAL RECEPTORS

- 1.1.1. The results of the modelling assessment at each modelled ecological receptor are presented in the below tables for each relevant pollutant and averaging period applicable to the study. The maximum modelled concentration and deposition values are presented, which is based on modelling over all five years of meteorological data (2016-2020). The change in PC and PEC, as a percentage of the relevant critical level / load, is presented for each receptor.
- 1.1.2. The PC impact in the with Proposed Scheme scenario represents the change in concentration / deposition between the Baseline scenario and Proposed Scheme scenario.
- 1.1.3. In all results tables presented in this appendix, the following designated site names are shortened to 'Thorne Moor SPA' and Thorne Moor SSSI', respectively:
 - a. Thorne & Hatfield Moors SPA
 - b. Thorne, Crowle and Goole Moors SSSI.
- 1.1.4. In the latest version of this Appendix, the results have been updated (see red table text) in response to changes to background pollutant concentrations and deposition on the APIS web platform in May 2023. Further details are provided in Air Quality Technical Note 3 (document reference 8.20) submitted at Deadline 8. The majority of changes on APIS do not affect the assessment outcome and, as such, updates in this Appendix have been limited to those that potentially impact on the significance of any effects. These include:
 - **a.** Reduction in background NOx concentrations over the Humber Estuary SAC/SPA/SSSI, where in comparison to data reported in Appendix 6.5 of the ES (APP-129), the site is moved from a situation where the PEC exceeds the critical level to a situation where the PEC is within the critical level, and
 - b. Increase in critical load for acid deposition over Lower Derwent Valley SAC and Barn Hill Meadows SSSI, where the site is moved from a situation where the PEC exceeds the critical load and the PC is greater than 1% of the critical load to a situation where the PEC is within the critical load and the maximum PC is less than 1% of the critical load.
- 1.1.4. In the latest version of this Appendix, the results reflect advice provided by Natural England via email that the May 2023 updates on the APIS website need not be incorporated into the assessment at this stage of the DCO process. Following this advice, the following change is made:
 - Background NOx concentrations over the Humber Estuary SAC/SPA/SSSI reflect
 APIS data from 2017-2019 within the 15km study area for the Proposed
 Development. In comparison to Revs 01 to 03 of this Appendix, this represents a

reduction in NOx concentrations since the original data presented reflected the maximum NOx concentrations *anywhere* within the Humber Estuary designated sites, and a slight increase in comparison to Rev 04 since this reflected the May 2023 updated (and reduced) 2019 – 2021 concentration data. Notwithstanding this update, as in Rev 04 of this Appendix, the NOx PEC over the Humber Estuary designated sites remains within the critical level both with and without the Proposed Development.

- 1.1.5. Further, the increased critical load for acid deposition over Lower Derwent Valley

 SAC and Barn Hill Meadows SSSI is retained from Rev 04 of this Appendix, and also applied to Breighton Meadows SSSI, Derwent Ings SSSI in this revision, This critical load was determined by the project ecologists and agreed with Natural England outwith the APIS update process. The rationale for this is presented in the Lower Derwent Valley Habitats and Soil Analysis Technical Note (REP3-009), which forms Appendix 8 to the HRA Report (REP6-021, Rev04 submitted at Deadline 9).
- 1.1.6. Finally the cumulative daily mean NOx process contributions (PCs) have been corrected in Table 1.14 for some designated sites where the annual mean NOx PCs had erroneously been inserted into the Table. This has no impact on the assessment since the amended PCs remain below the EA screening criterion (10% of the critical level) and the Predicted Environmental Concentrations (PECs), which did not require updating, are below the critical levels.

CORE MODEL SCENARIOS

1.1.5.1.1.7. Results pertaining to the core model scenarios are presented in **Tables 1.1 to**

Table 1.1 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NO_x

	Annual Mean NO _x concentration (μg/m³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	30	11.91	0.163	0.5%	12.13	40.4%		
Thorne Moor SAC/SPA/SSSI	30	13.21	0.076	0.3%	13.32	44.4%		
Lower Derwent SAC	30	9.92	0.166	0.6%	10.15	33.8%		
Lower Derwent SPA	30	9.92	0.166	0.6%	10.15	33.8%		
Skipwith Common SAC	30	9.76	0.058	0.2%	9.84	32.8%		
Skipwith Common SSSI	30	9.76	0.058	0.2%	9.84	32.8%		
Humber Estuary SAC	30	14.50 <u>19.22</u> ª	0.147	0.5%	<u>19.42</u> 14.70	<u>64.7</u> 49.0%		
Humber Estuary SPA/SSSI	30	14.50 <u>19.22</u> ^a	0.147	0.5%	<u>19.42</u> 14.70	<u>64.7</u> 49.0%		
Breighton Meadows SSSI	30	9.92	0.166	0.6%	10.15	33.8%		
Eskamhorn Meadows SSSI	30	11.35	0.046	0.2%	11.40	38.0%		
Derwent Ings SSSI	30	9.80	0.134	0.4%	9.99	33.3%		
Went Ings SSSI	30	12.09	0.052	0.2%	12.16	40.5%		
Barn Hill Meadows SSSI	30	12.89	0.152	0.5%	13.09	43.6%		
Burr Closes SSSI	30	10.53	0.062	0.2%	10.61	35.4%		
Common Plantation SINC	30	11.43	0.017	0.1%	11.45	38.2%		
Disused Railway Embankment SINC	30	10.76	0.040	0.1%	10.81	36.0%		
Barmby-on-the-Marsh LWS	30	10.48	0.076	0.3%	10.57	35.2%		
Brockholes SINC	30	11.22	0.019	0.1%	11.24	37.5%		
Meadow East of Orchard Farm SINC	30	10.83	0.009	0.0%	10.84	36.1%		
Barmby Pond LWS	30	9.96	0.127	0.4%	10.12	33.7%		
Cobble Croft Wood SINC	30	11.62	0.027	0.1%	11.65	38.8%		
Hagg Green Lane SINC	30	10.93	0.103	0.3%	11.07	36.9%		

		Annual Mean NO _x concentration (μg/m³)						
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Sand Pitt Wood & Barffs Close Plantation SINC	30	11.43	0.028	0.1%	11.46	38.2%		
Env. Agency Screening Criterion (as % of CL) 1% 70%								
a. Background NOx concentration for Humber Estuary SAC/SPA/S	SSI taken from maximum wi	thin study area (within 15km of Propos	sed Scheme, 2017-2019), from					

Table 1.2 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NO_x

Table 1.2 - Modelled Maximum Operational Impacts (Daily Mean NO _x concentration (μg/m³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	75	23.82	1.787	2.4%	27.96	37.3%		
Thorne Moor SAC/SPA/SSSI	75	26.42	0.893	1.2%	29.27	39.0%		
Lower Derwent SAC	75	19.84	0.893	1.2%	23.75	31.7%		
Lower Derwent SPA	75	19.84	0.893	1.2%	23.75	31.7%		
Skipwith Common SAC	75	19.52	0.646	0.9%	21.71	28.9%		
Skipwith Common SSSI	75	19.52	0.646	0.9%	21.71	28.9%		
Humber Estuary SAC	75	29.01 38.44 ^a	1.200	1.6%	<u>41.45</u> 32.02	<u>55.3</u> 42.7%		
Humber Estuary SPA/SSSI	75	29.01 38.44 ^a	1.200	1.6%	41.4532.02	<u>55.3</u> 42.7%		
Breighton Meadows SSSI	75	19.84	0.893	1.2%	23.75	31.7%		
Eskamhorn Meadows SSSI	75	22.70	1.912	2.5%	25.59	34.1%		
Derwent Ings SSSI	75	19.60	0.831	1.1%	23.37	31.2%		
Went Ings SSSI	75	24.18	0.971	1.3%	26.60	35.5%		
Barn Hill Meadows SSSI	75	25.78	0.962	1.3%	28.30	37.7%		
Burr Closes SSSI	75	21.06	0.761	1.0%	22.99	30.7%		
Common Plantation SINC	75	22.86	1.038	1.4%	24.19	32.3%		
Disused Railway Embankment SINC	75	21.52	1.184	1.6%	23.07	30.8%		

Receptor		Daily Mean NO _x concentration (μg/m³)							
	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Barmby-on-the-Marsh LWS	75	20.96	1.621	2.2%	23.44	31.3%			
Brockholes SINC	75	22.44	1.623	2.2%	24.44	32.6%			
Meadow East of Orchard Farm SINC	75	21.66	0.554	0.7%	22.33	29.8%			
Barmby Pond LWS	75	19.92	1.577	2.1%	23.29	31.1%			
Cobble Croft Wood SINC	75	23.24	1.176	1.6%	25.28	33.7%			
Hagg Green Lane SINC	75	21.86	1.248	1.7%	25.22	33.6%			
Sand Pitt Wood & Barffs Close Plantation SINC	75	22.86	1.630	2.2%	24.97	33.3%			
	- 1	Env. Agency Screening (Criterion (as % of CL)	10%					

a. Background NOx concentration for Humber Estuary SAC/SPA/SSSI taken from maximum within study area (within 15km of Proposed Scheme, 2017-2019), from

Table 1.3 - Modelled Maximum Operational Impacts at Ecological Receptors - Annual Mean NH₃

	Annual Mean NH₃ concentration (μg/m³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	3	4.57	0.007	0.2%	4.58	152.7%		
Thorne Moor SAC/SPA/SSSI	1	2.59	0.003	0.3%	2.60	259.5%		
Lower Derwent SAC	3	4.57	0.007	0.2%	4.58	152.7%		
Lower Derwent SPA	3	4.57	0.007	0.2%	4.58	152.7%		
Skipwith Common SAC	1	2.58	0.002	0.2%	2.58	258.4%		
Skipwith Common SSSI	1	2.58	0.002	0.2%	2.58	258.4%		
Humber Estuary SAC	3	3.58	0.004	0.1%	3.59	119.5%		
Humber Estuary SPA/SSSI	3	3.58	0.004	0.1%	3.59	119.5%		
Breighton Meadows SSSI	3	3.08	0.007	0.2%	3.09	103.0%		
Eskamhorn Meadows SSSI	3	2.40	0.002	0.1%	2.40	80.1%		

	Annual Mean NH₃ concentration (μg/m³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Derwent Ings SSSI	3	4.57	0.005	0.2%	4.58	152.6%		
Went Ings SSSI	3	2.35	0.002	0.1%	2.35	78.4%		
Barn Hill Meadows SSSI	3	2.32	0.005	0.2%	2.33	77.6%		
Burr Closes SSSI	3	2.50	0.003	0.1%	2.50	83.5%		
Common Plantation SINC	3	2.33	0.001	0.0%	2.33	77.7%		
Disused Railway Embankment SINC	1	2.28	0.002	0.1%	2.28	76.1%		
Barmby-on-the-Marsh LWS	3	2.28	0.003	0.1%	2.28	76.1%		
Brockholes SINC	3	2.28	0.001	0.0%	2.28	76.0%		
Meadow East of Orchard Farm SINC	1	2.33	0.000	0.0%	2.33	77.7%		
Barmby Pond LWS	1	2.28	0.006	0.2%	2.29	76.2%		
Cobble Croft Wood SINC	3	2.33	0.001	0.0%	2.33	77.7%		
Hagg Green Lane SINC	3	3.09	0.004	0.1%	3.10	103.2%		
Sand Pitt Wood & Barffs Close Plantation SINC	3	2.33	0.001	0.0%	2.33	77.7%		
		Env. Agency Screening C	Criterion (as % of CL)	1%		70%		

Table 1.4 - Modelled Maximum Operational Impacts at Ecological Receptors - Annual Mean SO₂

Receptor		Annual Mean SO ₂ concentration (μg/m³)							
	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
River Derwent SAC	20	3.93	0.072	0.4%	4.03	20.2%			
Thorne Moor SAC/SPA/SSSI	20	1.34	0.033	0.2%	1.39	7.0%			
Lower Derwent SAC	20	1.70	0.073	0.4%	1.81	9.0%			
Lower Derwent SPA	20	1.70	0.073	0.4%	1.81	9.0%			
Skipwith Common SAC	20	1.42	0.025	0.1%	1.46	7.3%			

		Annual Mean SO ₂ concentration (μg/m³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Skipwith Common SSSI	20	1.42	0.025	0.1%	1.46	7.3%			
Humber Estuary SAC	20	7.49	0.069	0.3%	7.59	38.0%			
Humber Estuary SPA/SSSI	20	7.49	0.069	0.3%	7.59	38.0%			
Breighton Meadows SSSI	20	1.70	0.073	0.4%	1.81	9.0%			
Eskamhorn Meadows SSSI	20	1.29	0.021	0.1%	1.32	6.6%			
Derwent Ings SSSI	20	1.69	0.059	0.3%	1.78	8.9%			
Went Ings SSSI	20	1.31	0.023	0.1%	1.34	6.7%			
Barn Hill Meadows SSSI	20	1.81	0.072	0.4%	1.91	9.5%			
Burr Closes SSSI	20	1.23	0.027	0.1%	1.27	6.3%			
Common Plantation SINC	20	1.44	0.008	0.0%	1.45	7.2%			
Disused Railway Embankment SINC	20	1.32	0.019	0.1%	1.34	6.7%			
Barmby-on-the-Marsh LWS	20	1.32	0.036	0.2%	1.36	6.8%			
Brockholes SINC	20	1.32	0.009	0.0%	1.33	6.7%			
Meadow East of Orchard Farm SINC	20	1.44	0.004	0.0%	1.44	7.2%			
Barmby Pond LWS	20	1.32	0.058	0.3%	1.40	7.0%			
Cobble Croft Wood SINC	20	1.44	0.013	0.1%	1.46	7.3%			
Hagg Green Lane SINC	20	1.43	0.047	0.2%	1.50	7.5%			
Sand Pitt Wood & Barffs Close Plantation SINC	20	1.44	0.013	0.1%	1.46	7.3%			
		Env. Agency Screening (Criterion (as % of CL)	1%		70%			

Table 1.5 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)								
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
River Derwent SAC	15	30.22	0.054	0.4%	30.30	202.0%			
Thorne Moor SAC	5	21.31	0.025	0.5%	21.35	426.9%			
Thorne Moor SPA	10	21.31	0.025	0.2%	21.35	213.5%			
Thorne Moor SSSI	5	21.31	0.025	0.5%	21.35	426.9%			
Lower Derwent SAC	20	30.22	0.055	0.3%	30.30	151.5%			
Lower Derwent SPA	20	30.22	0.055	0.3%	30.30	151.5%			
Skipwith Common SAC	10	21.12	0.019	0.2%	21.14	211.4%			
Skipwith Common SSSI	10	21.12	0.019	0.2%	21.14	211.4%			
Humber Estuary SAC	20	28.87	0.034	0.2%	28.92	144.6%			
Humber Estuary SPA/SSSI	20	28.87	0.034	0.2%	28.92	144.6%			
Breighton Meadows SSSI	20	23.51	0.055	0.3%	23.58	117.9%			
Eskamhorn Meadows SSSI	10	19.95	0.016	0.2%	19.97	199.7%			
Derwent Ings SSSI	20	30.22	0.043	0.2%	30.29	151.5%			
Went Ings SSSI	15	19.38	0.017	0.1%	19.41	129.4%			
Barn Hill Meadows SSSI	20	20.43	0.039	0.2%	20.48	102.4%			
Burr Closes SSSI	20	20.64	0.020	0.1%	20.67	103.3%			
Common Plantation SINC	10	33.74	0.010	0.1%	33.75	337.5%			
Disused Railway Embankment SINC	10	33.32	0.022	0.2%	33.35	333.5%			
Barmby-on-the-Marsh LWS	10	33.32	0.043	0.4%	33.37	333.7%			
Brockholes SINC	10	19.74	0.007	0.1%	19.75	197.5%			

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)								
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Meadow East of Orchard Farm SINC	20	19.88	0.003	0.0%	19.88	99.4%			
Barmby Pond LWS	10	19.74	0.044	0.4%	19.80	198.0%			
Cobble Croft Wood SINC	10	33.74	0.015	0.2%	33.76	337.6%			
Hagg Green Lane SINC	10	40.74	0.058	0.6%	40.82	408.2%			
Sand Pitt Wood & Barffs Close Plantation SINC	10	33.74	0.016	0.2%	33.76	337.6%			
Barlow Common LNR	10	33.74	0.010	0.1%	33.75	337.5%			
Env. Agency Screening C	riterion (as % of CL)			1%		70%			

Table 1.6 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Acid Deposition Rate

			Annual Acid Dep	osition Rate (keq/ha/yr))	
Receptor	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL
Thorne Moor SAC	0.462	1.73	0.006	1.3%	1.74	376.9%
Thorne Moor SSSI	0.462	1.73	0.006	1.3%	1.74	376.9%
Lower Derwent SAC	4.856	2.40	0.013	0.3%	2.42	49.9%
Skipwith Common SAC	0.802	1.73	0.005	0.6%	1.73	216.0%
Skipwith Common SSSI	0.802	1.73	0.005	0.6%	1.73	216.0%
Breighton Meadows SSSI	<u>4.856</u> 0.643	1.92	0.013	2.1% 0.3%	1.94	302.2% 40.0%
Eskamhorn Meadows SSSI	1.998	1.64	0.004	0.2%	1.64	82.2%
Derwent Ings SSSI	<u>4.856</u> 0.643	2.40	0.010	<u>0.2</u> 1.6%	2.42	376.5% 49.8%
Went Ings SSSI	2.008	1.59	0.004	0.2%	1.60	79.6%
Barn Hill Meadows SSSI	4.856	1.69	0.010	0.2%	1.70	35.1%
Burr Closes SSSI	1.248	1.68	0.005	0.4%	1.69	135.2%
	Env. Agency Screening Criterion (as % of CL)					70%

WITH PROPOSED SCHEME MITIGATION MODEL SCENARIOS

1.1.6.1.1.8. Results pertaining to the core model scenarios, including the With Proposed Scheme Mitigation (as detailed in **Section 6.10 of Chapter 6 (Air Quality)** (APP-042)) are presented in **Tables 1.7 to 1.12**.

Table 1.7 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NO_x (Including Mitigation)

	Annual Mean NO _x concentration (μg/m³)							
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation			
River Derwent SAC	30	0.163	0.133	0.5%	0.4%			
Thorne Moor SAC/SPA/SSSI	30	0.076	0.064	0.3%	0.2%			
Lower Derwent SAC	30	0.166	0.138	0.6%	0.5%			
Lower Derwent SPA	30	0.166	0.138	0.6%	0.5%			
Skipwith Common SAC	30	0.058	0.049	0.2%	0.2%			
Skipwith Common SSSI	30	0.058	0.049	0.2%	0.2%			
Humber Estuary SAC	30	0.147	0.125	0.5%	0.4%			
Humber Estuary SPA/SSSI	30	0.147	0.125	0.5%	0.4%			
Breighton Meadows SSSI	30	0.166	0.138	0.6%	0.5%			
Eskamhorn Meadows SSSI	30	0.046	0.034	0.2%	0.1%			
Derwent Ings SSSI	30	0.134	0.114	0.4%	0.4%			
Went Ings SSSI	30	0.052	0.042	0.2%	0.1%			
Barn Hill Meadows SSSI	30	0.152	0.124	0.5%	0.4%			
Burr Closes SSSI	30	0.062	0.050	0.2%	0.2%			
Common Plantation SINC	30	0.017	0.011	0.1%	0.0%			
Disused Railway Embankment SINC	30	0.040	0.029	0.1%	0.1%			
Barmby-on-the-Marsh LWS	30	0.076	0.057	0.3%	0.2%			
Brockholes SINC	30	0.019	0.013	0.1%	0.0%			
Meadow East of Orchard Farm SINC	30	0.009	0.006	0.0%	0.0%			
Barmby Pond LWS	30	0.127	0.098	0.4%	0.3%			
Cobble Croft Wood SINC	30	0.027	0.020	0.1%	0.1%			
Hagg Green Lane SINC	30	0.103	0.083	0.3%	0.3%			

Receptor		Annual Mean NO _x concentration (μg/m³)					
	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
Sand Pitt Wood & Barffs Close Plantation SINC	30	0.028	0.020	0.1%	0.1%		
Env. Agency Screening Criterion (as % of CL)					1%		

Table 1.8 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NO_x (Including Mitigation)

Table 1.8 - Modelled Maximum Operational Impact	Daily Mean NO _x concentration (μg/m³)							
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation			
River Derwent SAC	75	1.787	1.321	2.4%	1.8%			
Thorne Moor SAC/SPA/SSSI	75	0.893	0.699	1.2%	0.9%			
Lower Derwent SAC	75	0.893	0.673	1.2%	0.9%			
Lower Derwent SPA	75	0.893	0.673	1.2%	0.9%			
Skipwith Common SAC	75	0.646	0.434	0.9%	0.6%			
Skipwith Common SSSI	75	0.646	0.434	0.9%	0.6%			
Humber Estuary SAC	75	1.200	0.872	1.6%	1.2%			
Humber Estuary SPA/SSSI	75	1.200	0.872	1.6%	1.2%			
Breighton Meadows SSSI	75	0.893	0.673	1.2%	0.9%			
Eskamhorn Meadows SSSI	75	1.912	1.438	2.5%	1.9%			
Derwent Ings SSSI	75	0.831	0.670	1.1%	0.9%			
Went Ings SSSI	75	0.971	0.651	1.3%	0.9%			
Barn Hill Meadows SSSI	75	0.962	1.274	1.3%	1.7%			
Burr Closes SSSI	75	0.761	0.561	1.0%	0.7%			
Common Plantation SINC	75	1.038	0.684	1.4%	0.9%			
Disused Railway Embankment SINC	75	1.184	1.069	1.6%	1.4%			
Barmby-on-the-Marsh LWS	75	1.621	1.420	2.2%	1.9%			

Receptor	Daily Mean NO _x concentration (μg/m³)						
	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
Brockholes SINC	75	1.623	1.170	2.2%	1.6%		
Meadow East of Orchard Farm SINC	75	0.554	0.385	0.7%	0.5%		
Barmby Pond LWS	75	1.577	1.151	2.1%	1.5%		
Cobble Croft Wood SINC	75	1.176	0.767	1.6%	1.0%		
Hagg Green Lane SINC	75	1.248	0.832	1.7%	1.1%		
Sand Pitt Wood & Barffs Close Plantation SINC	75	1.630	1.083	2.2%	1.4%		
		Env. Agency S	1	10%			

Table 1.9 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NH₃ (Including Mitigation)

		Annual Mean NH ₃ concentration (μg/m³)							
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation				
River Derwent SAC	3	0.007	0.006	0.2%	0.2%				
Thorne Moor SAC/SPA/SSSI	1	0.003	0.003	0.3%	0.3%				
Lower Derwent SAC	3	0.007	0.006	0.2%	0.2%				
Lower Derwent SPA	3	0.007	0.006	0.2%	0.2%				
Skipwith Common SAC	1	0.002	0.002	0.2%	0.2%				
Skipwith Common SSSI	1	0.002	0.002	0.2%	0.2%				
Humber Estuary SAC	3	0.004	0.005	0.1%	0.2%				
Humber Estuary SPA/SSSI	3	0.004	0.005	0.1%	0.2%				
Breighton Meadows SSSI	3	0.007	0.006	0.2%	0.2%				
Eskamhorn Meadows SSSI	3	0.002	0.002	0.1%	0.1%				
Derwent Ings SSSI	3	0.005	0.005	0.2%	0.2%				
Went Ings SSSI	3	0.002	0.002	0.1%	0.1%				

			Annual Mean NH₃ concentrat	ion (µg/m³)	
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation
Barn Hill Meadows SSSI	3	0.005	0.005	0.2%	0.2%
Burr Closes SSSI	3	0.003	0.002	0.1%	0.1%
Common Plantation SINC	3	0.001	0.001	0.0%	0.0%
Disused Railway Embankment SINC	1	0.002	0.001	0.1%	0.0%
Barmby-on-the-Marsh LWS	3	0.003	0.003	0.1%	0.1%
Brockholes SINC	3	0.001	0.001	0.0%	0.0%
Meadow East of Orchard Farm SINC	1	0.000	0.000	0.0%	0.0%
Barmby Pond LWS	1	0.006	0.004	0.2%	0.1%
Cobble Croft Wood SINC	3	0.001	0.001	0.0%	0.0%
Hagg Green Lane SINC	3	0.004	0.004	0.1%	0.1%
Sand Pitt Wood & Barffs Close Plantation SINC	3	0.001	0.001	0.0%	0.0%
Env. Agency Screening Criterion (as % of CL)					1%

Table 1.10 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean SO₂ (Including Mitigation)

Receptor	Annual Mean SO₂ concentration (μg/m³)						
	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation		
River Derwent SAC	20	0.072	0.021	0.4%	0.1%		
Thorne Moor SAC/SPA/SSSI	20	0.033	0.008	0.2%	0.0%		
Lower Derwent SAC	20	0.073	0.020	0.4%	0.1%		
Lower Derwent SPA	20	0.073	0.020	0.4%	0.1%		
Skipwith Common SAC	20	0.025	0.007	0.1%	0.0%		
Skipwith Common SSSI	20	0.025	0.007	0.1%	0.0%		
Humber Estuary SAC	20	0.069	0.019	0.3%	0.1%		

		Annual Mean SO₂ concentration (μg/m³)							
Receptor	Critical Level	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation				
Humber Estuary SPA/SSSI	20	0.069	0.019	0.3%	0.1%				
Breighton Meadows SSSI	20	0.073	0.020	0.4%	0.1%				
Eskamhorn Meadows SSSI	20	0.021	0.006	0.1%	0.0%				
Derwent Ings SSSI	20	0.059	0.015	0.3%	0.1%				
Went Ings SSSI	20	0.023	0.007	0.1%	0.0%				
Barn Hill Meadows SSSI	20	0.072	0.020	0.4%	0.1%				
Burr Closes SSSI	20	0.027	0.008	0.1%	0.0%				
Common Plantation SINC	20	0.008	0.002	0.0%	0.0%				
Disused Railway Embankment SINC	20	0.019	0.005	0.1%	0.0%				
Barmby-on-the-Marsh LWS	20	0.036	0.011	0.2%	0.1%				
Brockholes SINC	20	0.009	0.003	0.0%	0.0%				
Meadow East of Orchard Farm SINC	20	0.004	0.001	0.0%	0.0%				
Barmby Pond LWS	20	0.058	0.017	0.3%	0.1%				
Cobble Croft Wood SINC	20	0.013	0.004	0.1%	0.0%				
Hagg Green Lane SINC	20	0.047	0.012	0.2%	0.1%				
Sand Pitt Wood & Barffs Close Plantation SINC	20	0.013	0.004	0.1%	0.0%				
		Env. Agency \$	Screening Criterion (as % of CL)						

Table 1.11 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate (Including Mitigation)

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)						
	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL - No Mitigation	Max PC Impact as % of CL - With Mitigation		
River Derwent SAC	15	0.054	0.044	0.4%	0.3%		
Thorne Moor SAC	5	0.025	0.021	0.5%	0.4%		

	Annual Nitrogen Deposition Rate (kgN/ha/yr)							
Receptor	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL - No Mitigation	Max PC Impact as % of CL - With Mitigation			
Thorne Moor SPA	10	0.025	0.021	0.2%	0.2%			
Thorne Moor SSSI	5	0.025	0.021	0.5%	0.4%			
Lower Derwent SAC	20	0.055	0.045	0.3%	0.2%			
Lower Derwent SPA	20	0.055	0.045	0.3%	0.2%			
Skipwith Common SAC	10	0.019	0.016	0.2%	0.2%			
Skipwith Common SSSI	10	0.019	0.016	0.2%	0.2%			
Humber Estuary SAC	20	0.034	0.041	0.2%	0.2%			
Humber Estuary SPA/SSSI	20	0.034	0.041	0.2%	0.2%			
Breighton Meadows SSSI	20	0.055	0.045	0.3%	0.2%			
Eskamhorn Meadows SSSI	10	0.016	0.012	0.2%	0.1%			
Derwent Ings SSSI	20	0.043	0.037	0.2%	0.2%			
Went Ings SSSI	15	0.017	0.014	0.1%	0.1%			
Barn Hill Meadows SSSI	20	0.039	0.041	0.2%	0.2%			
Burr Closes SSSI	20	0.020	0.016	0.1%	0.1%			
Common Plantation SINC	10	0.010	0.007	0.1%	0.1%			
Disused Railway Embankment SINC	10	0.022	0.016	0.2%	0.2%			
Barmby-on-the-Marsh LWS	10	0.043	0.032	0.4%	0.3%			
Brockholes SINC	10	0.007	0.005	0.1%	0.0%			
Meadow East of Orchard Farm SINC	20	0.003	0.002	0.0%	0.0%			
Barmby Pond LWS	10	0.044	0.033	0.4%	0.3%			
Cobble Croft Wood SINC	10	0.015	0.011	0.2%	0.1%			
Hagg Green Lane SINC	10	0.058	0.046	0.6%	0.5%			

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)						
	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL - No Mitigation	Max PC Impact as % of CL – With Mitigation		
Sand Pitt Wood & Barffs Close Plantation SINC	10	0.016	0.011	0.2%	0.1%		
Barlow Common LNR	10	0.010	0.007	0.1%	0.1%		
	Env. Agency Screening Criterion (as % of CL)						

Table 1.12 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Acid Deposition Rate (Including Mitigation)

			Annual Acid Deposition Rate	(keq/ha/yr)		
Receptor	Critical Load	Max PC Impact – No Mitigation	Max PC Impact – With Mitigation	Max PC Impact as % of CL – No Mitigation	Max PC Impact as % of CL – With Mitigation	
Thorne Moor SAC	0.462	0.006	0.003	1.3%	0.6%	
Thorne Moor SSSI	0.462	0.006	0.003	1.3%	0.6%	
Lower Derwent SAC	4.856	0.013	0.006	0.3%	0.1%	
Skipwith Common SAC	0.802	0.005	0.002	0.6%	0.3%	
Skipwith Common SSSI	0.802	0.005	0.002	0.6%	0.3%	
Breighton Meadows SSSI	4.8560.643	0.013	0.006	2.1% 0.3%	<u>0.1</u> 1.0%	
Eskamhorn Meadows SSSI	1.998	0.004	0.002	0.2%	0.1%	
Derwent Ings SSSI	4.8560.643	0.010	0.005	<u>0.2</u> 1.6%	0. <u>1</u> 8%	
Went Ings SSSI	2.008	0.004	0.002	0.2%	0.1%	
Barn Hill Meadows SSSI	4.856	0.010	0.006	0.2%	0.1%	
Burr Closes SSSI	1.248	0.005	0.002	0.4%	0.2%	
		Env. Agency \$	Screening Criterion (as % of CL)	1%		

CUMULATIVE IMPACTS (WITH PROPOSED SCHEME & OTHER PROJECTS)

1.1.7.1.1.9. Results pertaining to the cumulative impacts, both before mitigation ("No Mit") and after mitigation ("Mitig") associated With Proposed Scheme Mitigation is applied (as detailed in **Section 6.10 of Chapter 6 (Air Quality)**) are presented in **Tables 1.13 to 1.18**.

Table 1.13 - Modelled Maximum Cumulative Impacts at Ecological Receptors - Annual Mean NO_x

	Annual Mean NO _x concentration (μg/m³)											
Receptor	Critical Level	Max Cumulat	ive PC Impact	Max PC Impac	t as % of CL	Max Cumu	lative PEC	Max PEC a	s % of CL			
	Citical Level	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ			
River Derwent SAC	30	0.675	0.647	2.3%	2.2%	12.63	12.60	42.1%	42.0%			
Thorne Moor SAC/SPA/SSSI	30	0.348	0.336	1.2%	1.1%	13.58	13.57	45.3%	45.2%			
Lower Derwent SAC	30	0.681	0.653	2.3%	2.2%	10.64	10.62	35.5%	35.4%			
Lower Derwent SPA	30	0.681	0.653	2.3%	2.2%	10.64	10.62	35.5%	35.4%			
Skipwith Common SAC	30	0.504	0.495	1.7%	1.7%	10.29	10.28	34.3%	34.3%			
Skipwith Common SSSI	30	0.504	0.495	1.7%	1.7%	10.29	10.28	34.3%	34.3%			
Humber Estuary SAC	30	0.498	0.474	1.7%	1.6%	<u>19.77</u> 15.05	<u>19.74</u> 15.03	<u>65.9</u> 50.2 %	<u>65.8</u> 50.1%			
Humber Estuary SPA/SSSI	30	0.498	0.474	1.7%	1.6%	<u>19.77</u> 15.05	<u>19.74</u> 15.03	<u>65.9</u> 50.2%	<u>65.8</u> 50.1%			
Breighton Meadows SSSI	30	0.681	0.653	2.3%	2.2%	10.64	10.62	35.5%	35.4%			
Eskamhorn Meadows SSSI	30	0.488	0.483	1.6%	1.6%	11.84	11.83	39.5%	39.4%			
Derwent Ings SSSI	30	0.640	0.619	2.1%	2.1%	10.49	10.47	35.0%	34.9%			
Went Ings SSSI	30	0.303	0.296	1.0%	1.0%	12.40	12.40	41.3%	41.3%			
Barn Hill Meadows SSSI	30	0.557	0.527	1.9%	1.8%	13.48	13.45	44.9%	44.8%			
Burr Closes SSSI	30	0.283	0.276	0.9%	0.9%	10.83	10.82	36.1%	36.1%			
Common Plantation SINC	30	0.728	0.723	2.4%	2.4%	12.16	12.15	40.5%	40.5%			
Disused Railway Embankment SINC	30	0.558	0.546	1.9%	1.8%	11.32	11.31	37.7%	37.7%			
Barmby-on-the-Marsh LWS	30	0.575	0.555	1.9%	1.8%	11.07	11.05	36.9%	36.8%			
Brockholes SINC	30	0.480	0.476	1.6%	1.6%	11.70	11.70	39.0%	39.0%			
Meadow East of Orchard Farm SINC	30	0.701	0.699	2.3%	2.3%	11.53	11.53	38.4%	38.4%			
Barmby Pond LWS	30	0.611	0.585	2.0%	2.0%	10.59	10.57	35.3%	35.2%			
Cobble Croft Wood SINC	30	0.697	0.690	2.3%	2.3%	12.32	12.31	41.1%	41.0%			
Hagg Green Lane SINC	30	0.704	0.684	2.3%	2.3%	11.67	11.65	38.9%	38.8%			

		Annual Mean NO _x concentration (μg/m³)										
Receptor	Critical Level	Max Cumulative PC Impact		Max PC Impact as % of CL		Max Cumulative PEC		Max PEC as % of CL				
		No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ			
Sand Pitt Wood & Barffs Close Plantation SINC	30	0.798	0.790	2.7%	2.6%	12.23	12.22	40.8%	40.7%			
Env. Agency Screening	Env. Agency Screening Criterion (as % of CL)							70%	%			

Table 1.14 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Daily Mean NO_x

Table 1.14 - Modelled Maximum Cumulative impa	Daily Mean NO _x concentration (μg/m³)												
Receptor	Critical Level	Max Cumula	tive PC Impact	Max PC Impa	ct as % of CL	Max Cumul	ative PEC	Max PEC	as % of CL				
	Critical Level	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ				
River Derwent SAC	75	5.459	5.078	7.3%	6.8%	31.44	31.24	41.9%	41.7%				
Thorne Moor SAC/SPA/SSSI	75	<u>4.336</u> 0.446	4.1410.446	<u>5.8%</u> 0.6%	<u>5.5%</u> 0.6%	32.43	32.34	43.2%	43.1%				
Lower Derwent SAC	75	4.3820.308	4.1990.308	<u>5.8%</u> 0.4%	<u>5.6%</u> 0.4%	27.00	26.90	36.0%	35.9%				
Lower Derwent SPA	75	4.3820.308	4.1990.308	<u>5.8%</u> 0.4%	<u>5.6%</u> 0.4%	27.00	26.90	36.0%	35.9%				
Skipwith Common SAC	75	<u>3.932</u> 0.253	<u>3.615</u> 0.253	<u>5.2%</u> 0.3%	4.8%0.3%	24.75	24.66	33.0%	32.9%				
Skipwith Common SSSI	75	3.932 0.253	<u>3.615</u> 0.253	<u>5.2%</u> 0.3%	4.8%0.3%	24.75	24.66	33.0%	32.9%				
Humber Estuary SAC	75	4.6090.469	4.3650.469	<u>6.1%</u> 0.6%	<u>5.8%</u> 0.6%	44.5735.14	44.4535.01	<u>59.4%</u> 46.9%	<u>59.3%</u> 46.7%				
Humber Estuary SPA/SSSI	75	4.6090.469	4.3650.469	<u>6.1%</u> 0.6%	<u>5.8%</u> 0.6%	44.5735.14	44.4535.01	<u>59.4%</u> 46.9%	<u>59.3%</u> 46.7%				
Breighton Meadows SSSI	75	<u>4.382</u> 0.306	<u>4.198</u> 0.306	<u>5.8%</u> 0.4%	<u>5.6%</u> 0.4%	27.00	26.89	36.0%	35.9%				
Eskamhorn Meadows SSSI	75	<u>6.777</u> 0.983	6.3470.983	<u>9.0%</u> 1.3%	<u>8.5%</u> 1.3%	30.40	29.99	40.5%	40.0%				
Derwent Ings SSSI	75	<u>4.234</u> 0.308	<u>4.158</u> 0.308	<u>5.6%</u> 0.4%	<u>5.5%</u> 0.4%	26.63	26.47	35.5%	35.3%				
Went Ings SSSI	75	<u>5.111</u> 0.245	5.0400.245	<u>6.8%</u> 0.3%	<u>6.7%</u> 0.3%	30.27	30.14	40.4%	40.2%				
Barn Hill Meadows SSSI	75	<u>4.591</u> 0.691	4.4780.691	<u>6.1%</u> 0.9%	<u>6.0%</u> 0.9%	32.25	32.14	43.0%	42.9%				
Burr Closes SSSI	75	3.7360.346	3.5370.346	<u>5.0%</u> 0.5%	<u>4.7%</u> 0.5%	25.99	25.95	34.7%	34.6%				
Common Plantation SINC	75	7.240	7.596	9.7%	10.1%	30.30	30.65	40.4%	40.9%				
Disused Railway Embankment SINC	75	6.035	5.779	8.0%	7.7%	27.93	27.62	37.2%	36.8%				

		Daily Mean NO _x concentration (μg/m³)											
Receptor	Cuitical Laval	Max Cumulat	tive PC Impact	Max PC Impac	ct as % of CL	Max Cumulative PEC		Max PEC as % of CL					
	Critical Level	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ				
Barmby-on-the-Marsh LWS	75	6.310	5.938	8.4%	7.9%	28.32	27.95	37.8%	37.3%				
Brockholes SINC	75	5.892	5.440	7.9%	7.3%	28.71	28.26	38.3%	37.7%				
Meadow East of Orchard Farm SINC	75	5.436	5.702	7.2%	7.6%	27.18	27.44	36.2%	36.6%				
Barmby Pond LWS	75	5.189	4.763	6.9%	6.4%	27.29	27.09	36.4%	36.1%				
Cobble Croft Wood SINC	75	7.028	6.706	9.4%	8.9%	31.22	30.90	41.6%	41.2%				
Hagg Green Lane SINC	75	5.261	4.843	7.0%	6.5%	29.27	28.95	39.0%	38.6%				
Sand Pitt Wood & Barffs Close Plantation SINC	75	7.720	7.948	10.3%	10.6%	31.06	31.02	41.4%	41.4%				
	Env. Agency Screening Criterion (as % of CL)			109	%								

Table 1.15 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean NH₃

				Annual Mean	NH ₃ concentra	ation (µg/m³)			
Receptor	Critical Level	Max Cumulat	ive PC Impact	Max PC Impac	ct as % of CL	Max Cumul	ative PEC	Max PEC a	as % of CL
	Critical Level	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ
River Derwent SAC	3	0.008	0.007	0.3%	0.2%	4.58	4.58	152.7%	152.7%
Thorne Moor SAC/SPA/SSSI	1	0.006	0.006	0.6%	0.6%	2.60	2.60	259.8%	259.8%
Lower Derwent SAC	3	0.008	0.007	0.3%	0.2%	4.58	4.58	152.7%	152.7%
Lower Derwent SPA	3	0.008	0.007	0.3%	0.2%	4.58	4.58	152.7%	152.7%
Skipwith Common SAC	1	0.003	0.003	0.3%	0.3%	2.58	2.58	258.4%	258.4%
Skipwith Common SSSI	1	0.003	0.003	0.3%	0.3%	2.58	2.58	258.4%	258.4%
Humber Estuary SAC	3	0.006	0.007	0.2%	0.2%	3.59	3.59	119.6%	119.7%
Humber Estuary SPA/SSSI	3	0.006	0.007	0.2%	0.2%	3.59	3.59	119.6%	119.7%
Breighton Meadows SSSI	3	0.008	0.007	0.3%	0.2%	3.09	3.09	103.0%	103.0%
Eskamhorn Meadows SSSI	3	0.003	0.003	0.1%	0.1%	2.40	2.40	80.1%	80.1%

				Annual Mean	NH₃ concentra	ation (µg/m³)			
Receptor	Critical Level	Max Cumulat	ive PC Impact	Max PC Impac	ct as % of CL	Max Cumul	ative PEC	Max PEC	as % of CL
	Childai Levei	No Mit ⁿ	Mitig ⁿ						
Derwent Ings SSSI	3	0.006	0.005	0.2%	0.2%	4.58	4.58	152.7%	152.6%
Went Ings SSSI	3	0.004	0.004	0.1%	0.1%	2.36	2.35	78.5%	78.5%
Barn Hill Meadows SSSI	3	0.006	0.007	0.2%	0.2%	2.33	2.33	77.6%	77.7%
Burr Closes SSSI	3	0.003	0.003	0.1%	0.1%	2.50	2.50	83.5%	83.5%
Common Plantation SINC	3	0.002	0.001	0.1%	0.0%	2.33	2.33	77.7%	77.7%
Disused Railway Embankment SINC	3	0.003	0.002	0.1%	0.1%	2.28	2.28	76.1%	76.1%
Barmby-on-the-Marsh LWS	3	0.004	0.004	0.1%	0.1%	2.29	2.28	76.2%	76.1%
Brockholes SINC	3	0.002	0.002	0.1%	0.1%	2.28	2.28	76.1%	76.1%
Meadow East of Orchard Farm SINC	3	0.001	0.001	0.0%	0.0%	2.33	2.33	77.7%	77.7%
Barmby Pond LWS	3	0.007	0.005	0.2%	0.2%	2.29	2.29	76.3%	76.2%
Cobble Croft Wood SINC	3	0.002	0.002	0.1%	0.1%	2.33	2.33	77.7%	77.7%
Hagg Green Lane SINC	3	0.005	0.004	0.2%	0.1%	3.10	3.10	103.2%	103.2%
Sand Pitt Wood & Barffs Close Plantation SINC	3	0.002	0.002	0.1%	0.1%	2.33	2.33	77.7%	77.7%
	Env. Agency Screening Criterion (as % of CL)			19	/ 6			70%	

Table 1.16 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Mean SO₂

	Annual Mean SO₂ concentration (μg/m³)											
Receptor	Critical Level	Max Cumulat	ive PC Impact	Max PC Impac	t as % of CL	Max Cumul	ative PEC	Max PEC as % of CL				
	Critical Level	No Mit ⁿ	Mitig ⁿ									
River Derwent SAC	20	0.074	0.022	0.4%	0.1%	4.04	3.98	20.2%	19.9%			
Thorne Moor SAC/SPA/SSSI	20	0.041	0.017	0.2%	0.1%	1.40	1.38	7.0%	6.9%			
Lower Derwent SAC	20	0.074	0.022	0.4%	0.1%	1.81	1.76	9.0%	8.8%			
Lower Derwent SPA	20	0.074	0.022	0.4%	0.1%	1.81	1.76	9.0%	8.8%			

	Annual Mean SO₂ concentration (μg/m³)												
Receptor	Critical Lavel	Max Cumulat	ive PC Impact	Max PC Impac	t as % of CL	Max Cumul	ative PEC	Max PEC as	% of CL				
	Critical Level	No Mit ⁿ	Mitig ⁿ										
Skipwith Common SAC	20	0.026	0.008	0.1%	0.0%	1.46	1.44	7.3%	7.2%				
Skipwith Common SSSI	20	0.026	0.008	0.1%	0.0%	1.46	1.44	7.3%	7.2%				
Humber Estuary SAC	20	0.073	0.023	0.4%	0.1%	7.59	7.54	38.0%	37.7%				
Humber Estuary SPA/SSSI	20	0.073	0.023	0.4%	0.1%	7.59	7.54	38.0%	37.7%				
Breighton Meadows SSSI	20	0.074	0.022	0.4%	0.1%	1.81	1.76	9.0%	8.8%				
Eskamhorn Meadows SSSI	20	0.024	0.009	0.1%	0.0%	1.32	1.30	6.6%	6.5%				
Derwent Ings SSSI	20	0.061	0.017	0.3%	0.1%	1.78	1.74	8.9%	8.7%				
Went Ings SSSI	20	0.028	0.011	0.1%	0.1%	1.35	1.33	6.7%	6.7%				
Barn Hill Meadows SSSI	20	0.075	0.024	0.4%	0.1%	1.91	1.86	9.6%	9.3%				
Burr Closes SSSI	20	0.028	0.009	0.1%	0.0%	1.27	1.25	6.3%	6.3%				
Common Plantation SINC	20	0.009	0.004	0.0%	0.0%	1.45	1.44	7.3%	7.2%				
Disused Railway Embankment SINC	20	0.021	0.008	0.1%	0.0%	1.34	1.33	6.7%	6.7%				
Barmby-on-the-Marsh LWS	20	0.038	0.013	0.2%	0.1%	1.37	1.34	6.8%	6.7%				
Brockholes SINC	20	0.011	0.005	0.1%	0.0%	1.33	1.33	6.7%	6.6%				
Meadow East of Orchard Farm SINC	20	0.006	0.003	0.0%	0.0%	1.45	1.44	7.2%	7.2%				
Barmby Pond LWS	20	0.060	0.019	0.3%	0.1%	1.40	1.36	7.0%	6.8%				
Cobble Croft Wood SINC	20	0.014	0.005	0.1%	0.0%	1.46	1.45	7.3%	7.2%				
Hagg Green Lane SINC	20	0.049	0.014	0.2%	0.1%	1.50	1.47	7.5%	7.3%				
Sand Pitt Wood & Barffs Close Plantation SINC	20	0.014	0.005	0.1%	0.0%	1.46	1.45	7.3%	7.2%				
	Env. Agency Screening Criterion (as % of C				6		70%						

Table 1.17 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate

		,	- Allitual Nitrogel	Annual Nitroge		Rate (kgN/ha/yr))		
Receptor	Critical Load	Max Cumula	ative PC Impact	Max PC Impa	ct as % of CL	Max Cumu	lative PEC	Max PEC	as % of CL
	Critical Load	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ
River Derwent SAC	15	0.106	0.098	0.7%	0.7%	30.35	30.34	202.3%	202.3%
Thorne Moor SAC	5	0.061	0.063	1.2%	1.3%	21.38	21.38	427.6%	427.6%
Thorne Moor SPA	10	0.061	0.063	0.6%	0.6%	21.38	21.38	213.8%	213.8%
Thorne Moor SSSI	5	0.061	0.063	1.2%	1.3%	21.38	21.38	427.6%	427.6%
Lower Derwent SAC	20	0.105	0.098	0.5%	0.5%	30.35	30.34	151.8%	151.7%
Lower Derwent SPA	20	0.105	0.098	0.5%	0.5%	30.35	30.34	151.8%	151.7%
Skipwith Common SAC	10	0.067	0.064	0.7%	0.6%	21.19	21.19	211.9%	211.9%
Skipwith Common SSSI	10	0.067	0.064	0.7%	0.6%	21.19	21.19	211.9%	211.9%
Humber Estuary SAC	20	0.073	0.086	0.4%	0.4%	28.96	28.98	144.8%	144.9%
Humber Estuary SPA/SSSI	20	0.073	0.086	0.4%	0.4%	28.96	28.98	144.8%	144.9%
Breighton Meadows SSSI	20	0.105	0.098	0.5%	0.5%	23.63	23.62	118.2%	118.1%
Eskamhorn Meadows SSSI	10	0.058	0.061	0.6%	0.6%	20.01	20.02	200.1%	200.2%
Derwent Ings SSSI	20	0.096	0.091	0.5%	0.5%	30.34	30.34	151.7%	151.7%
Went Ings SSSI	15	0.049	0.046	0.3%	0.3%	19.44	19.43	129.6%	129.6%
Barn Hill Meadows SSSI	20	0.082	0.088	0.4%	0.4%	20.52	20.54	102.6%	102.7%
Burr Closes SSSI	20	0.045	0.041	0.2%	0.2%	20.69	20.69	103.5%	103.4%
Common Plantation SINC	10	0.158	0.155	1.6%	1.5%	33.90	33.90	339.0%	339.0%
Disused Railway Embankment SINC	10	0.126	0.129	1.3%	1.3%	33.45	33.45	334.5%	334.5%
Barmby-on-the-Marsh LWS	10	0.141	0.141	1.4%	1.4%	33.47	33.47	334.7%	334.7%
Brockholes SINC	10	0.056	0.057	0.6%	0.6%	19.80	19.80	198.0%	198.0%
Meadow East of Orchard Farm SINC	20	0.075	0.076	0.4%	0.4%	19.96	19.96	99.8%	99.8%
Barmby Pond LWS	10	0.095	0.085	1.0%	0.9%	19.85	19.84	198.5%	198.4%

				Annual Nitroge	n Deposition F	Rate (kgN/ha/yr)				
Receptor	Critical Load	Max Cumulative PC Impact		Max PC Impa	ct as % of CL	Max Cumulative PEC		Max PEC as % of CL		
	Critical Load	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	
Cobble Croft Wood SINC	10	0.155	0.151	1.6%	1.5%	33.90	33.89	339.0%	338.9%	
Hagg Green Lane SINC	10	0.176	0.173	1.8%	1.7%	40.94	40.93	409.4%	409.3%	
Sand Pitt Wood & Barffs Close Plantation SINC	10	0.176	0.171	1.8%	1.7%	33.92	33.91	339.2%	339.1%	
Barlow Common LNR	10	0.173	0.170	1.7%	1.7%	33.91	33.91	339.1%	339.1%	
Env. Agency Screening Criterion (as % of CL)				1%				70%		

Table 1.18 - Modelled Maximum Cumulative Impacts at Ecological Receptors – Annual Acid Deposition Rate

		-		Annual Acid [Deposition Rat	e (keq/ha/yr)			
Receptor	Critical Load	Max Cumulat	ive PC Impact	Max PC Impa	ct as % of CL	Max Cumul	ative PEC	Max PEC	as % of CL
	Critical Load	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ	No Mit ⁿ	Mitig ⁿ
Thorne Moor SAC	0.4623	0.010	0.007	2.1%	1.5%	1.74	1.74	375.8%	377.1%
Thorne Moor SSSI	0.4623	0.010	0.007	2.1%	1.5%	1.75	1.74	377.8%	377.1%
Lower Derwent SAC	4.856	0.017	0.010	0.4%	0.2%	2.43	2.42	50.0%	49.8%
Skipwith Common SAC	0.802	0.008	0.006	1.0%	0.7%	1.74	1.73	216.4%	216.1%
Skipwith Common SSSI	0.802	0.008	0.006	1.0%	0.7%	1.74	1.73	216.4%	216.1%
Breighton Meadows SSSI	0.643	0.017	0.010	<u>0.4</u> 2.6%	0.21.6%	1.95	1.94	302.8 <u>40.1</u> %	<u>40.0</u> 301.7%
Eskamhorn Meadows SSSI	1.998	0.007	0.005	0.3%	0.3%	1.64	1.64	82.3%	82.2%
Derwent Ings SSSI	0.643	0.014	0.009	<u>0.3</u> 2.2%	0.21.4%	2.42	2.42	<u>49.9</u> 377.1%	<u>49.8</u> 376.2%
Went Ings SSSI	2.008	0.007	0.005	0.4%	0.2%	1.60	1.60	79.7%	79.6%
Barn Hill Meadows SSSI	4.856	0.014	0.010	0.3%	0.2%	1.71	1.70	35.2%	35.1%
Burr Closes SSSI	1.248	0.007	0.004	0.5%	0.3%	1.69	1.69	135.3%	135.1%
	Env. Agency Screening Criterion (as % of CL)			19	%			70%	

SENSITIVITY TEST: WORST CASE EMISSIONS PROFILE

- 1.1.8.1.1.10. Results pertaining to the worst-case emissions profile sensitivity test are presented in **Tables 1.19 to 1.24**, based on emissions from the Proposed Scheme alone.
- 4.1.9.1.1.1. For all pollutant concentrations and deposition rates, it is evident that the modelled maximum PC impacts attributed to the with Proposed Scheme scenario are lower at all receptors relative to the core model scenarios. This is a function of all four biomass units in the Baseline scenario switching from 'midmerit' operation (full load for 4,000 hours per year) to continuous operation (full load for 8,760 hours per year), resulting in more pollutants being emitted and thus more pronounced changes (increases) in concentrations / deposition rates relative to the With Proposed Scheme scenario. In the With Proposed Scheme scenario, operation changes from 'mid-merit' to continuous full load at the two non-BECCS biomass units only (BECCS units already assumed to operate at continuous full load in core modelling scenario), meaning the changes (increases and decreases) in concentrations / deposition rates are relatively small compared to the Baseline.
- 4.1.10.1.1.12. As a consequence, the maximum modelled impacts of the Proposed Scheme decrease at all receptors under the worst-case emissions profile scenario relative to the core modelling. Whilst some modelled maximum PEC concentrations do increase under worst case emissions in both the Baseline and With Proposed Scheme scenarios, there are no material changes relative to the core modelling equivalents, meaning that the respective assessment significance criteria are not exceeded.
- 1.1.11.13. The results confirm that the assessment of likely significant effects reported in **Chapter 6 (Air Quality)** is not affected when considering the worst-case emissions profiles in both the Baseline and With Proposed Scheme scenarios. Given that the modelled maximum impacts are lower under a worst-case emissions profile, there was no need to repeat the test in relation to cumulative impacts, as the core modelling results for the cumulative scenarios represent the most conservative results in terms of potential impacts.

Table 1.19 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean NO_x (Worst Case Emissions Profile)

	Annual Mean NO _x concentration (μg/m³)								
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
River Derwent SAC	30	11.91	0.055	0.2%	12.10	40.3%			
Thorne Moor SAC/SPA/SSSI	30	13.21	0.026	0.1%	13.33	44.4%			
Lower Derwent SAC	30	9.92	0.058	0.2%	10.11	33.7%			
Lower Derwent SPA	30	9.92	0.058	0.2%	10.11	33.7%			
Skipwith Common SAC	30	9.76	0.023	0.1%	9.84	32.8%			
Skipwith Common SSSI	30	9.76	0.023	0.1%	9.84	32.8%			
Humber Estuary SAC	30	<u>19.22</u> 14.50	0.056	0.2%	<u>19.40</u> 14.68	<u>64.7</u> 48.9%			
Humber Estuary SPA/SSSI	30	<u>19.22</u> 14.50	0.056	0.2%	<u>19.40</u> 14.68	<u>64.7</u> 48.9%			
Breighton Meadows SSSI	30	9.92	0.058	0.2%	10.11	33.7%			
Eskamhorn Meadows SSSI	30	11.35	0.012	0.0%	11.38	37.9%			
Derwent Ings SSSI	30	9.80	0.048	0.2%	9.98	33.3%			
Went Ings SSSI	30	12.09	0.018	0.1%	12.15	40.5%			
Barn Hill Meadows SSSI	30	12.89	0.055	0.2%	13.04	43.5%			
Burr Closes SSSI	30	10.53	0.020	0.1%	10.59	35.3%			
Common Plantation SINC	30	11.43	0.004	0.0%	11.44	38.1%			
Disused Railway Embankment SINC	30	10.76	0.009	0.0%	10.78	35.9%			
Barmby-on-the-Marsh LWS	30	10.48	0.021	0.1%	10.53	35.1%			
Brockholes SINC	30	11.22	0.004	0.0%	11.23	37.4%			
Meadow East of Orchard Farm SINC	30	10.83	0.002	0.0%	10.83	36.1%			
Barmby Pond LWS	30	9.96	0.036	0.1%	10.06	33.5%			
Cobble Croft Wood SINC	30	11.62	0.007	0.0%	11.64	38.8%			
Hagg Green Lane SINC	30	10.93	0.034	0.1%	11.05	36.8%			

	Annual Mean NO _x concentration (μg/m³)							
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
Sand Pitt Wood & Barffs Close Plantation SINC	30	30 11.43 0.007			11.45	38.2%		
		Env. Agency Screening C	riterion (as % of CL)	1%		70%		

Table 1.20 - Modelled Maximum Operational Impacts at Ecological Receptors – Daily Mean NO_x (Worst Case Emissions Profile)

			Daily Mean NO	x concentration (μg/m³)		
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL
River Derwent SAC	75	23.82	1.787	2.4%	27.96	37.3%
Thorne Moor SAC/SPA/SSSI	75	26.42	0.893	1.2%	29.27	39.0%
Lower Derwent SAC	75	19.84	0.893	1.2%	23.75	31.7%
Lower Derwent SPA	75	19.84	0.893	1.2%	23.75	31.7%
Skipwith Common SAC	75	19.52	0.646	0.9%	21.71	28.9%
Skipwith Common SSSI	75	19.52	0.646	0.9%	21.71	28.9%
Humber Estuary SAC	75	38.4429.01	1.200	1.6%	41.4532.02	<u>55.3</u> 42.7%
Humber Estuary SPA/SSSI	75	<u>38.44</u> 29.01	1.200	1.6%	<u>41.45</u> 32.02	<u>55.3</u> 42.7%
Breighton Meadows SSSI	75	19.84	0.893	1.2%	23.75	31.7%
Eskamhorn Meadows SSSI	75	22.70	1.912	2.5%	25.59	34.1%
Derwent Ings SSSI	75	19.60	0.831	1.1%	23.37	31.2%
Went Ings SSSI	75	24.18	0.971	1.3%	26.60	35.5%
Barn Hill Meadows SSSI	75	25.78	0.962	1.3%	28.30	37.7%
Burr Closes SSSI	75	21.06	0.761	1.0%	22.99	30.7%
Common Plantation SINC	75	22.86	1.038	1.4%	24.19	32.3%
Disused Railway Embankment SINC	75	21.52	1.184	1.6%	23.07	30.8%
Barmby-on-the-Marsh LWS	75	20.96	1.621	2.2%	23.44	31.3%

	Daily Mean NO _x concentration (μg/m³)								
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Brockholes SINC	75	22.44	1.623	2.2%	24.44	32.6%			
Meadow East of Orchard Farm SINC	75	21.66	0.554	0.7%	22.33	29.8%			
Barmby Pond LWS	75	19.92	1.577	2.1%	23.29	31.1%			
Cobble Croft Wood SINC	75	23.24	1.176	1.6%	25.28	33.7%			
Hagg Green Lane SINC	75	21.86	1.248	1.7%	25.22	33.6%			
Sand Pitt Wood & Barffs Close Plantation SINC	75	22.86	1.630	2.2%	24.97	33.3%			
	10%								

Table 1.21 - Modelled Maximum Operational Impacts at Ecological Receptors - Annual Mean NH₃ (Worst Case Emissions Profile)

			Annual Mean NH	₃ concentration (μg/m³)		
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL
River Derwent SAC	3	4.57	0.002	0.1%	4.58	152.6%
Thorne Moor SAC/SPA/SSSI	1	2.59	0.001	0.1%	2.60	259.6%
Lower Derwent SAC	3	4.57	0.002	0.1%	4.58	152.6%
Lower Derwent SPA	3	4.57	0.002	0.1%	4.58	152.6%
Skipwith Common SAC	1	2.58	0.001	0.1%	2.58	258.4%
Skipwith Common SSSI	1	2.58	0.001	0.1%	2.58	258.4%
Humber Estuary SAC	3	3.58	0.002	0.1%	3.59	119.6%
Humber Estuary SPA/SSSI	3	3.58	0.002	0.1%	3.59	119.6%
Breighton Meadows SSSI	3	3.08	0.002	0.1%	3.09	103.0%
Eskamhorn Meadows SSSI	3	2.40	0.001	0.0%	2.40	80.1%
Derwent Ings SSSI	3	4.57	0.002	0.1%	4.58	152.6%
Went Ings SSSI	3	2.35	0.001	0.0%	2.35	78.4%

			Annual Mean NH	3 concentration (μg/m³)		
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL
Barn Hill Meadows SSSI	3	2.32	0.002	0.1%	2.33	77.6%
Burr Closes SSSI	3	2.50	0.001	0.0%	2.50	83.4%
Common Plantation SINC	3	2.33	0.000	0.0%	2.33	77.7%
Disused Railway Embankment SINC	1	2.28	0.000	0.0%	2.28	76.0%
Barmby-on-the-Marsh LWS	3	2.28	0.001	0.0%	2.28	76.1%
Brockholes SINC	3	2.28	0.000	0.0%	2.28	76.0%
Meadow East of Orchard Farm SINC	1	2.33	0.000	0.0%	2.33	77.7%
Barmby Pond LWS	1	2.28	0.001	0.0%	2.29	76.2%
Cobble Croft Wood SINC	3	2.33	0.000	0.0%	2.33	77.7%
Hagg Green Lane SINC	3	3.09	0.001	0.0%	3.10	103.2%
Sand Pitt Wood & Barffs Close Plantation SINC	3	2.33	0.000	0.0%	2.33	77.7%
	1%		70%			

Table 1.22 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Mean SO₂ (Worst Case Emissions Profile)

	Annual Mean SO₂ concentration (μg/m³)								
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
River Derwent SAC	20	3.93	0.021	0.1%	4.03	20.1%			
Thorne Moor SAC/SPA/SSSI	20	1.34	0.009	0.0%	1.40	7.0%			
Lower Derwent SAC	20	1.70	0.021	0.1%	1.80	9.0%			
Lower Derwent SPA	20	1.70	0.021	0.1%	1.80	9.0%			
Skipwith Common SAC	20	1.42	0.008	0.0%	1.46	7.3%			
Skipwith Common SSSI	20	1.42	0.008	0.0%	1.46	7.3%			
Humber Estuary SAC	20	7.49	0.021	0.1%	7.58	37.9%			

			Annual Mean SO	2 concentration (μg/m³)		
Receptor	Critical Level	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL
Humber Estuary SPA/SSSI	20	7.49	0.021	0.1%	7.58	37.9%
Breighton Meadows SSSI	20	1.70	0.021	0.1%	1.80	9.0%
Eskamhorn Meadows SSSI	20	1.29	0.005	0.0%	1.31	6.5%
Derwent Ings SSSI	20	1.69	0.016	0.1%	1.78	8.9%
Went Ings SSSI	20	1.31	0.007	0.0%	1.34	6.7%
Barn Hill Meadows SSSI	20	1.81	0.022	0.1%	1.89	9.4%
Burr Closes SSSI	20	1.23	0.008	0.0%	1.26	6.3%
Common Plantation SINC	20	1.44	0.002	0.0%	1.44	7.2%
Disused Railway Embankment SINC	20	1.32	0.004	0.0%	1.33	6.7%
Barmby-on-the-Marsh LWS	20	1.32	0.009	0.0%	1.35	6.7%
Brockholes SINC	20	1.32	0.002	0.0%	1.32	6.6%
Meadow East of Orchard Farm SINC	20	1.44	0.001	0.0%	1.44	7.2%
Barmby Pond LWS	20	1.32	0.015	0.1%	1.37	6.9%
Cobble Croft Wood SINC	20	1.44	0.003	0.0%	1.45	7.2%
Hagg Green Lane SINC	20	1.43	0.013	0.1%	1.49	7.5%
Sand Pitt Wood & Barffs Close Plantation SINC	20	1.44	0.003	0.0%	1.45	7.2%
		Env. Agency Screening C	Criterion (as % of CL)	1%		70%

Table 1.23 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Nitrogen Deposition Rate (Worst Case Emissions Profile)

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)							
Receptor	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL		
River Derwent SAC	15	30.22	0.017	0.1%	30.29	201.9%		
Thorne Moor SAC	5	21.31	0.008	0.2%	21.35	427.0%		

Becomton	Annual Nitrogen Deposition Rate (kgN/ha/yr)									
Receptor	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL				
Thorne Moor SPA	10	21.31	0.008	0.1%	21.35	213.5%				
Thorne Moor SSSI	5	21.31	0.008	0.2%	21.35	427.0%				
Lower Derwent SAC	20	30.22	0.017	0.1%	30.29	151.5%				
Lower Derwent SPA	20	30.22	0.017	0.1%	30.29	151.5%				
Skipwith Common SAC	10	21.12	0.007	0.1%	21.14	211.4%				
Skipwith Common SSSI	10	21.12	0.007	0.1%	21.14	211.4%				
Humber Estuary SAC	20	28.87	0.017	0.1%	28.93	144.7%				
Humber Estuary SPA/SSSI	20	28.87	0.017	0.1%	28.93	144.7%				
Breighton Meadows SSSI	20	23.51	0.017	0.1%	23.57	117.9%				
Eskamhorn Meadows SSSI	10	19.95	0.004	0.0%	19.96	199.6%				
Derwent Ings SSSI	20	30.22	0.014	0.1%	30.29	151.4%				
Went Ings SSSI	15	19.38	0.005	0.0%	19.40	129.4%				
Barn Hill Meadows SSSI	20	20.43	0.017	0.1%	20.49	102.4%				
Burr Closes SSSI	20	20.64	0.006	0.0%	20.66	103.3%				
Common Plantation SINC	10	33.74	0.002	0.0%	33.75	337.5%				
Disused Railway Embankment SINC	10	33.32	0.005	0.1%	33.33	333.3%				
Barmby-on-the-Marsh LWS	10	33.32	0.011	0.1%	33.35	333.5%				
Brockholes SINC	10	19.74	0.001	0.0%	19.74	197.4%				
Meadow East of Orchard Farm SINC	20	19.88	0.001	0.0%	19.88	99.4%				
Barmby Pond LWS	10	19.74	0.012	0.1%	19.78	197.8%				
Cobble Croft Wood SINC	10	33.74	0.004	0.0%	33.75	337.5%				
Hagg Green Lane SINC	10	40.74	0.017	0.2%	40.81	408.1%				
Sand Pitt Wood & Barffs Close Plantation SINC	10	33.74	0.004	0.0%	33.75	337.5%				

Receptor	Annual Nitrogen Deposition Rate (kgN/ha/yr)								
	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL			
Barlow Common LNR	10 33.74 0.002 0.0		0.0%	33.75	337.5%				
	Env. Age	ncy Screening Cr	iterion (as % of CL)	1%		70%			

Table 1.24 - Modelled Maximum Operational Impacts at Ecological Receptors – Annual Acid Deposition Rate (Worst Case Emissions Profile)

			Annual Acid Dep	osition Rate (keq/ha/yr))	
Receptor	Critical Load	Background	Max PC Impact	Max PC Impact as % of CL	Proposed Scheme Max PEC	Max PEC as % of CL
Thorne Moor SAC	0.462	1.73	0.002	0.4%	1.74	377.2%
Thorne Moor SSSI	0.462	1.73	0.002	0.4%	1.74	377.2%
Lower Derwent SAC	4.856	2.40	0.004	0.1%	2.42	49.9%
Skipwith Common SAC	0.802	1.73	0.002	0.2%	1.73	216.0%
Skipwith Common SSSI	0.802	1.73	0.002	0.2%	1.73	216.0%
Breighton Meadows SSSI	4.8560.643	1.92	0.004	0. <u>1</u> 6%	1.94	<u>40.0</u> 301.9%
Eskamhorn Meadows SSSI	1.998	1.64	0.001	0.0%	1.64	82.1%
Derwent Ings SSSI	4.8560.643	2.40	0.003	0. <u>1</u> 5%	2.42	<u>49.8</u> 376.4%
Went Ings SSSI	2.008	1.59	0.001	0.1%	1.60	79.5%
Barn Hill Meadows SSSI	4.856	1.69	0.004	0.1%	1.70	35.1%
Burr Closes SSSI	1.248	1.68	0.001	0.1%	1.69	135.1%
	1%		70%			